

# HARD X-RAY AND GAMMA-RAY OBSERVATIONS OF SN1987A DURING THE PERIOD 1987-1989

S. M. Schindler, W. R. Cook, D. M. Palmer, T. A. Prince, C. H. Starr,  
and E. C. Stone

California Institute of Technology, Pasadena, California 91125 USA

## Abstract

The Caltech Gamma-Ray Imaging Payload (GRIP) has observed SN1987A on four balloon flights from Alice Springs, Australia, in May and Nov. 1987, April 1988, and April 1989 (days: 85, 269, 414, and 771 respectively). The instrument is a coded-aperture telescope, sensitive to radiation in the energy range 30 keV to 10 MeV (Althouse *et al.* 1985). Both gamma-ray continuum and line emission were detected from the supernova, in qualitative agreement with existing models. In addition, we have obtained the first images of the SN1987A region at gamma-ray energies, confirming that the bulk of the gamma-ray emission comes from the supernova, and not from LMC X-1 (Cook *et al.* 1988). We review the results of these observations, including flux levels and upper limits for both line emission and Compton scattered continuum for each of the four flights.

This work supported in part by NASA grant NGR 05-002-160

## References

- Althouse W.E. *et al.* 1985 19th ICRC Conference Papers **3**, 299-302  
Cook W.R. *et al.* 1988 Ap. J.(*Letters*) **334**, L87-L90